

ON AFFINE ALGEBRAS

Martin Lorenz
Max-Planck-Institut für Mathematik
Gottfried-Claren-Str. 26
D-5300 Bonn 3, Fed. Rep. Germany

These notes contain a unified approach, via bimodules, to a number of results of Artin-Tate type. Throughout we will keep the following notation:

k is a commutative ring (with 1), and
 R and S are k -algebras.

As is customary and convenient, (R,S) -bimodules V are assumed to have identical k -operations on both sides: $\xi v = v\xi$ ($v \in V, \xi \in k$).

1. BIMODULES AND AFFINE ALGEBRAS

LEMMA 1. Let $0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$ be a short exact sequence of (R,S) -bimodules and assume that V_S and ${}_R W$ are finitely generated, say $V = Rv_1 + \dots + Rv_m + U$ for suitable $v_i \in V$.

If S is affine over k , then there exists an affine k -subalgebra $R' \subseteq R$ and a finitely generated (R',S) -subbimodule $U' \subseteq U$ such that

$$V = R'v_1 + \dots + R'v_m + U' .$$

PROOF. Write $V = w_1 S + \dots + w_n S$ and let $x_1, \dots, x_t \in S$ be k -algebra generators for S . Then

$$w_i = \sum_{h=1}^m r_{ih} v_h + u_i, \quad v_i x_j = \sum_{h=1}^m r_{ijh} v_h + u_{ij}$$

for suitable $r_{ih}, r_{ijh} \in R$ and $u_i, u_{ij} \in U$. Let $R' \subseteq R$ be the k -subalgebra generated by the r_{ih} 's and r_{ijh} 's, and let $U' \subseteq U$ be the (R',S) -subbimodule generated by the u_i 's and u_{ij} 's. Then $V' = R'v_1 + \dots + R'v_m + U'$ contains w_i and $v_h x_j$ for all i, j, h .

Hence

$$V'x_j = \sum_{h=1}^m R'v_h x_j + U'x_j \subseteq R'V' + U' = V' .$$

Since $V'k = kV' \subseteq V'$, it follows that $\sum_{i=1}^n w_i S \subseteq V'S \subseteq V'$, whence $V' = V$.

■

COROLLARY 1. Let $R \subseteq S$ be k -algebras such that S is affine over k and finitely generated as a left module over R . Then S is also finitely generated as a left module over some affine subalgebra $R' \subseteq R$, with the same module generators.

PROOF. Take $U = 0$ and $V = {}_R S_S$ in the lemma.

■

Recall that, for a left R -module V , the trace of V in R is defined by

$$\text{Tr}_R(V) = \sum \{ \text{Im } f \mid f \in \text{Hom}_R(V, R) \} .$$

$\text{Tr}_R(V)$ is a two-sided ideal of R , and V is a generator for $R\text{-mod}$, the category of left R -modules, if and only if $\text{Tr}_R(V) = R$.

LEMMA 2. Let V be an (R, S) -bimodule such that ${}_R V$ and V_S are finitely generated, and assume that S is affine over k . Suppose that R contains an affine k -subalgebra $A \subseteq R$ and a finitely generated left ideal I with $I \subseteq \text{Tr}_R(V)$ and $R = \langle A, I \rangle_{k\text{-algebra}}$ ($= A + IA$) . Then R is affine over k . This happens in particular if ${}_R V$ is a generator for $R\text{-mod}$.

PROOF. By assumption on I , there exist finitely many $f_i \in \text{Hom}_R(V, R)$ with $I \subseteq \sum \text{Im } f_i$. After enlarging I if necessary we may therefore assume that a finite direct sum of copies of ${}_R V$ maps onto I . By Lemma 1, with $U = 0$, there exists an affine k -subalgebra

$R' \subseteq R$ such that ${}_R V$ is finitely generated. Hence ${}_R I$ is also finitely generated, and A, R' , and the generators of I over R' together generate $\langle A, I \rangle_{k\text{-algebra}} = R$.

COROLLARY 2. Let $R \subseteq S$ be k -algebras with S affine over k . Assume that S and $\text{Tr}_R(S)$ are finitely generated as left modules over R . Then R is affine over k if and only if $R/\text{Tr}_R(S)$ is affine over k .

PROOF. Apply Lemma 2 with $V = {}_R S_S$ and $I = \text{Tr}_R(S)$.

2. SOME APPLICATIONS

(A) CORNERS OF RINGS. Assume that S is affine over k and let $e = e^2 \in S$. If eS is finitely generated as left ideal of S , then eSe is affine over k . (Montgomery-Small [6]).

PROOF. By [6, Lemma 1], eS is finitely generated as left module over eSe . Now take $V = eS$ and $R = eSe$ in Lemma 2 and note that ${}_R V$ maps onto ${}_R R$ via $es \mapsto ese (S \in S)$.

(B) MORITA EQUIVALENCE. If A and B are Morita equivalent rings, then there exists an (A, B) -bimodule P such that ${}_A P$ and P_B are finitely generated projective generators for $A\text{-mod}$, resp. $\text{mod-}B$. In case A and B are k -algebras, and the left and right k -operations on P agree, we conclude from Lemma 2 that A is affine over k if and only if B is affine over k . (Wadsworth, cf. [6, Acknowledgement]).

(C) RESULTS OF ARTIN-TATE TYPE. Let $R \subseteq S$ be k -algebras with S affine over k and ${}_R S$ finitely generated. Then R is affine over k in each of the following cases:

- i. R is a finitely generated left module over a commutative subalgebra and k is Noetherian;
- ii. S is left Noetherian and $R \subseteq S$ is a finite centralizing extension (i.e., $S = \sum_{i=1}^n Rx_i$ with $x_i r = r s_i$ for all $r \in R$);
- iii. ${}_R S$ is projective and, for each proper two-sided ideal M of R , $MS \neq S$ (e.g., if ${}_R S$ is free or if ${}_R S$ is projective and maximal ideals of R are localizable);
- iv. k is Noetherian and, for some commutative subalgebra $C \subseteq R$, the module $\left(\frac{S}{R}\right)_C$ is Noetherian.

PROOF. (i). One can clearly assume that R itself is commutative. Choose $R' \subseteq R$ as in Corollary 1. Then R' is Noetherian, by the Hilbert basis theorem, and hence ${}_R R'$ is finitely generated, as ${}_R S$ is. Thus R is affine.

(ii). Again, Corollary 1 yields $R' \subseteq R$ affine such that $R' \subseteq S$ is a finite centralizing extension. As S is left Noetherian, the Eisenbud-Eakin theorem [3] implies that R' is likewise. Now argue as in (i).

(iii). Set $T = \text{Tr}_R(S)$. Then $TS = S$, by the dual basis lemma, and so we must have $T = R$. (Actually, by [2], ${}_R S$ maps onto ${}_R R$.) The result now follows from Corollary 2.

(iv). Let $C \subseteq R$ be commutative with $\left(\frac{S}{R}\right)_C$ Noetherian and set $X = \{r \in R \mid Sr \subseteq R\} = \text{ann}\left(\frac{S}{R}\right)_R$. Then $X = SX \subseteq \text{Tr}_R({}_R S)$ and $X = \bigcap_v \text{rt. ann}_R(v+R)$, where v runs over a finite generating set for $\left(\frac{S}{R}\right)_R$. Therefore, $\left(\frac{R}{X}\right)_R \hookrightarrow \left(\frac{S}{R}\right)_R^n$ for some n . Since $\left(\frac{S}{R}\right)_C$ is Noetherian, we conclude that $\left(\frac{R}{X}\right)_C$ is Noetherian, and hence $\left(\frac{S}{X}\right)_C$ is also Noetherian. By Lemma 1, with $V = {}_S S_C$ and $W = \left(\frac{S}{X}\right)_C$ (and with sides interchanged), we can find an affine subalgebra $C' \subseteq C$ and a finitely generated (S, C') -subbimodule $X' \subseteq X$ with $\left(\frac{S}{X'}\right)_C$ finitely generated. Now C' is Noetherian and so $\left(\frac{R}{X'}\right)_C$ is finitely generated too. Moreover, since ${}_R S$ is finitely generated, X' is also finitely generated as (R, C') -bimodule, say

$$X' = \sum_{i=1}^n R x_i C' .$$

Now set $I = \sum_{i=1}^n R x_i$, so that I is a finitely generated left ideal of R with $I \subseteq \text{Tr}_R(S)$, and let $A \subseteq R$ be the subalgebra generated by C' and the generators of $\left(\frac{R}{X'}\right)_{C'}$. Then A is affine and $R = A + X' = A + IA$. Thus Lemma 2 yields the result. ■

REMARKS. (i) is a mild generalization of the original Artin-Tate Lemma [1] and has been observed by a number of people.

(ii) is contained in [6]. Using a result of Formanek and Jategaonkar [4] instead of the Eisenbud-Eakin theorem, the same proof yields versions of (ii) which work for certain finite normalizing extensions $R \subseteq S$. For example, if $S = \sum_{i=1}^n R x_i$ with $rx_i = x_i r^{\sigma_i}$ for certain automorphisms σ_i of R and if $G = \langle \sigma_1, \dots, \sigma_n \rangle$ acts locally finitely on R , then the argument goes through, because we can then choose $R' \subseteq R$ to be affine and normalized by x_i 's. Also, for any finite normalizing extension $R \subseteq S$, proper right ideals of R extend to proper right ideals of S [5]. Thus (iii) above applies to finite normalizing extensions $R \subseteq S$ with ${}_R S$ projective. The question as to whether the Artin-Tate lemma holds for general finite normalizing extensions $R \subseteq S$, with S left Noetherian, say, was raised in [6] and is still open as far as I know.

For the moment, let T denote the class of finitely generated left R -modules V such that $\text{Tr}_R(V)V = V$. Then the assumptions in (iii) could be replaced by: ${}_R S \in T$ and, for each proper two-sided ideal M of R , $MS \neq S$. Now T contains all finitely generated projective modules over R as well as, clearly, all generators of $R\text{-mod}$, and T is closed under direct sums. But I don't know of an easy characterization of the modules in T .

The prototype of (iv) (with $C = k$) is due to Lance Small (oral communication).

ACKNOWLEDGEMENT. Research supported by the Deutsche Forschungsgemeinschaft/Heisenberg Programm (Lo 261/2-2). I would like to thank Lance Small for numerous interesting conversations about affine algebras and other things.

REFERENCES.

- [1] E. Artin and J.T. Tate: A note on finite ring extensions, *J. Math. Soc. Japan* 3 (1951), 74-77.
- [2] B. Cortzen, L.W. Small and J.T. Stafford: Decomposing overrings, *Proc. Amer. Math. Soc.* 82 (1981), 28-30.
- [3] D. Eisenbud: Subrings of Artinian and Noetherian rings, *Math. Ann.* 185 (1970), 247-249.
- [4] E. Formanek and A.V. Jategaonkar: Subrings of Noetherian rings, *Proc. Amer. Math. Soc.* 46 (1974), 181-186.
- [5] M. Lorenz: Finite normalizing extensions of rings, *Math. Z.* 176 (1981), 447-484.
- [6] S. Montgomery and L.W. Small: Fixed rings of Noetherian rings, *Bull. London Math. Soc.* 13 (1981), 33-38.